This listing of claims will replace all prior versions, and listings of claims in the application.

## In the Claims:

- (currently amended) A method for detecting an isolate of a B. cereus group, in a sample, the method comprising:
  - (a) placing on a mieroehip microarray of claim 10 oligonucleotide probes targeted to rRNA sequences wherein at least one mismatch is sufficient to discriminate among the B. cereus subgroups;
  - providing conditions for hybridization of the probes with rRNA from the sample; and
  - analyzing hybridization signals in the microehip microarray from which the particular isolate is detected.
- (currently amended) The method of claim 1, wherein the oligonucleotide probes
  are directed to 16S rRNA and 23S rRNA. and the corresponding target sequences are shown in
  FIG. 1 and FIG. 2.
  - 3. (original) The method of claim 1, wherein the probes are labeled.
- (original) The method of claim 3, wherein the labels are selected from the group consisting of fluorescent dyes, radio isotopes, immunological labels, immuno-chemical labels and gold particles.
- (currently amended) The method of claim 1, wherein the oligonucleotide probes selected from the group consisting of sequences listed in Table 5 discriminate one or more subgroups Anthracis, Cereus A, Cereus B, Thuringiensis A, Thuringiensis B, Mycoides A and Mycoides B.
- (Currently amended) The method of claim 1, wherein pairs of oligonucleotide probes that discriminate subgroups Anthracis from Cereus A are ps21/ps22, ps23/ps24, 23F13/23F14, 23F15/23F16, and C9/C10.
- (original) The method of claim 1, wherein a ratio of hybridization signals of oligonucleotide probes ps17 and ps18 discriminates between B. anthracis Ames and B. cereus 9620.

- (currently amended) The method of claim 1, wherein the oligonucleotide probes are ps21; and ps22; ps23; and ps24 to discriminate B. anthracis Sterne from B. cereus HER 1414 and B. thuringenesis B8.
- (withdrawn) The method of claim 1, wherein the oligonucleotide probes are ps7, ps8, and ps9 to discriminate B. thuringenesis 4Q281 from other B. cereus subgroup isolates.
- (currently amended) A microarray that comprises oligonucleotide probes selected from the group consisting of sequences designated:

Oligonucleotide Name	5' to 3' Sequence
ps5	CCG CTA ACT TCA TAA GAG CA (SEQ ID NO: 74)
ps6	CCG CTA ACT TCt TgA GAG CA (SEQ ID NO: 75)
ps17	TCT AGG GTT GTC AGA GGA TG (SEQ ID NO: 86)
ps18	TCT AGG GTT tTC AGA GGA TG (SEQ ID NO: 87)
ps19	TCT GCT CCC GAA GGA GAA GC (SEQ ID NO: 88)
ps20	TCT GCc CCC GAA GGg GAA GC (SEQ ID NO: 89)
ps21	CAG CTC AGC CTT CAC GAT AA (SEQ ID NO: 90)
ps22	CAG CTC AGC CTT tAC GAT AA (SEQ ID NO: 91)
23F1	TTT GGG CTA TGT TCC GTT TC (SEQ ID NO: 126) and
23F2	TTT GGG CTA GAT TCC GTT TC (SEQ ID NO: 127)
<u>A8</u>	CCC TCT GTG ACG AGC CTT TC (SEQ ID NO: 143)

11. (previously presented) The microarray of claim 10, wherein the oligonucleotides are arranged in a specific pattern wherein I, II, III and IV are columns and A, B, C, D, E, and F are rows in the microarray:

	I	II	III	IV
A	ps19	ps20		
В				
C			ps5	ps6
D			=	-
E				
F			ps17	ps18

12. (previously presented) A microarray as in claim 10, wherein the oligonucleotides are arranged in pairs; ps19 and ps20; ps5 and ps6; ps17 and ps18.

- 13. (canceled).
- 14. (canceled).
- (canceled).
- 16. (currently amended) A probe selected from the group consisting of sequences designated:

Oligonucleotide Name	5' to 3' Sequence
ps5	CCG CTA ACT TCA TAA GAG CA (SEQ ID NO: 74)
ps6	CCG CTA ACT TCt TgA GAG CA (SEQ ID NO: 75)
ps17	TCT AGG GTT GTC AGA GGA TG (SEQ ID NO: 86)
ps18	TCT AGG GTT tTC AGA GGA TG (SEQ ID NO: 87)
ps19	TCT GCT CCC GAA GGA GAA GC (SEQ ID NO: 88)
ps20	TCT GCc CCC GAA GGg GAA GC (SEQ ID NO: 89)
ps21	CAG CTC AGC CTT CAC GAT AA (SEQ ID NO: 90)
ps22	CAG CTC AGC CTT tAC GAT AA (SEQ ID NO: 91)
	TTT GGG CTA TGT TCC GTT TC (SEQ ID NO: 126) and
	TTT GGG CTA GAT TCC GTT TC (SEQ ID NO: 127) and
<u>A8</u>	CCC TCT GTG ACG AGC CTT TC (SEQ ID NO: 143).

- 17. (previously presented) The probe of claim 16, wherein the sequence of the oligonucleotide probe is the full reverse complement of a sequence of claim 16.
  - 18. (canceled).
  - 19. (canceled).
- 20. (currently amended) A diagnostic kit to detect B. anthracis target rRNA in a sample, the diagnostic kit comprising:
  - (a) a microchip that comprises at least one oligonucleotide probe that distinguishes B. anthracis from other closely related microorganisms, wherein the oligonucleotide is selected from the group consisting of

CCG CTA ACT TCA TAA GAG CA (SEQ ID NO: 74), CCG CTA ACT TCt TgA GAG CA (SEQ ID NO: 75), TCT AGG GTT GTC AGA GGA TG (SEQ ID NO: 86), TCT AGG GTT tTC AGA GGA TG (SEQ ID NO: 87), TCT GCT CCC GAA GGA GAA GC (SEQ ID NO: 88), TCT GCc CCC GAA GGg GAA GC (SEQ ID NO: 89), CAG CTC AGC CTT CAC GAT AA (SEQ ID NO: 90), CAG CTC AGC CTT tAC GAT AA (SEQ ID NO: 91), TTT GGG CTA TGT TCC GTT TC (SEQ ID NO: 126), TTT GGG CTA GAT TCC GTT TC (SEQ ID NO: 127), and CCC TCT GTG ACG AGC CTT TC (SEQ ID NO: 143); and

- (b) a method for detecting hybridization between the at least one probe and the target rRNA by which hybridization, B. anthracis is detected.
- (withdrawn) A method for taxonomically classifying B. cereus groups, said method comprising:
  - developing strain- and subgroup-specific signature profiles of 16S and 23S rRNA sequences for B. cereus group isolates; and
  - (b) using the signature profiles to construct phylogenetic trees in order to classify the various B. cereus group isolates.
  - 22. (canceled).